

ODD-SYMPLECTIC GROUP IN FIRST ORDER PARTIAL DIFFERENTIAL EQUATIONS

L.SBANO

ABSTRACT. In this paper it is shown that the characteristic vector field associated to a first order PDE:

$$h(x, \nabla z(x)) = 0 \text{ with } x \in \mathbb{R}^n$$

has the same form of an infinitesimal generator of an odd-symplectic transformation with contact Hamiltonian $h(x, p)$ on the level set $h = 0$. We also study under which condition such PDE has a characteristic vector field commuting with a generator of an odd-symplectic action.

INTRODUCTION

Very recently R. Cushman [4] presented the notion of *odd-symplectic group*. This notion was formulated in [6] but differently in [7]; more recently in [5, 3]. Such group can be defined on vector spaces with odd dimension. Loosely speaking, the odd-symplectic group lies between the standard symplectic group and the contact group. More precisely, it interpolates symplectic groups in different dimensions according to the following exact sequence:

$$\dots \rightarrow Sp(2n+2, \mathbb{R}) \rightarrow Sp(2n+1, \mathbb{R}) \rightarrow Sp(2n, \mathbb{R}) \rightarrow \dots$$

Although the definition of the odd-symplectic group is natural, at the moment, there are few examples of applications. The aim of this paper is to show that odd-symplectic transformations rise naturally in the geometric theory of first order PDE and in particular in the study of the characteristics of PDEs of the following form:

$$(1) \quad h(x, \nabla z(x)) = 0,$$

where $x \in \mathbb{R}^n$ $h(.,.)$ is a smooth function on $\mathbb{R}^{2n}$.

Here is an outline of the paper: in the first section the odd-symplectic group is defined. In the following sections we recall the theory of characteristics for first order PDEs and we show that the characteristic vector field associated to (1) coincides with an infinitesimal generator of an odd-symplectic transformation. We show also that

$$h(x, \nabla z(x)) = \frac{1}{2} \langle g \nabla z, \nabla z \rangle + \langle v, \nabla z \rangle + k$$

with g constant $n \times n$ symmetric matrix, $k \in \mathbb{R}$ and $v \in \mathbb{R}^n$ has a symmetry whose infinitesimal generator is given by an element in the Lie algebra of $Sp(2n+1, \mathbb{R})$. Finally we show that there is an isomorphism between certain class of first order PDE and the Lie algebra of the odd-symplectic group; moreover we show the connection of the odd-symplectic group with Hamilton-Jacobi theory and the Eikonal equation.

1. ODD-SYMPLECTIC GROUP AND CONTACT GEOMETRY

Let us consider a symplectic vector space (V, ω) , $\dim V = 2n + 2$. The symplectic group is:

$$(2) \quad Sp(V, \omega) \doteq \{A \in Gl(V) | A^* \omega = \omega\}.$$

The odd-symplectic group $Sp(V, \omega)_{v_0}$ can be defined as follows: let $v_0 \in V$ be a non-zero vector, then:

$$(3) \quad Sp(V, \omega)_{v_0} \doteq \{A \in Sp(V, \omega) | Av_0 = v_0\}$$

Sometimes $Sp(V, \omega)_{v_0}$ is denoted by $Sp(2n + 1, \mathbb{R})$. This group has been studied in [3, 5]. It turns out that the odd symplectic group arises naturally in contact geometry. Let us follow [3]. Consider a contact manifold $(\mathcal{M}, \theta)$ ($\dim \mathcal{M} = 2n + 1$) where locally, in Darboux coordinates, the contact form θ is:

$$(4) \quad \theta = dz - y_i dx^i$$

In [1] it is shown that one can symplectify $(\mathcal{M}, \theta)$ into $(\mathcal{N}, \omega) = (\mathbb{R}_+ \times M, d(t\theta))$ where $t > 0$ is the coordinate on $\mathbb{R}_+$. Consider a one parameter group of symplectomorphisms $F_s : \mathcal{N} \rightarrow \mathcal{N}$ of the form:

$$(5) \quad F_s(t, m) = (t, G_s(t, m)).$$

As shown in [3] the infinitesimal generator of F_s at $t = 1, y = 0$ is an element of the algebra $sp(2n + 1, \mathbb{R})$ of $Sp(2n + 1, \mathbb{R})$. If X is vector field on $\mathcal{N}$ such that

$$(6) \quad L_X(t) = 0 \text{ and } L_X(d(t\theta)) = 0$$

then its flow is a map of the form (5). Since

$$L_X(d(t\theta)) = d(i_X(d(t\theta))) = 0$$

we deduce that (6) implies:

$$(7) \quad L_X(t) = 0 \text{ and } i_X d(t\theta) = \widehat{dh} \text{ for some } \widehat{h} : \mathcal{N} \rightarrow \mathbb{R}$$

To see what this means, let us use Darboux coordinates. If we denote

$$X_{\widehat{h}} = \frac{dt}{ds} \frac{\partial}{\partial t} + \frac{dz}{ds} \frac{\partial}{\partial z} + \frac{dx^i}{ds} \frac{\partial}{\partial x^i} + \frac{dy_i}{ds} \frac{\partial}{\partial y_i}$$

then (7) gives:

$$(8) \quad \begin{cases} \frac{dt}{ds} = 0 \\ \frac{dx^i}{ds} = \frac{1}{t} \frac{\partial \widehat{h}}{\partial y_i} \\ \frac{dy_i}{ds} = -\frac{1}{t} \frac{\partial \widehat{h}}{\partial x^i} \\ \frac{dz}{ds} = -\frac{\partial \widehat{h}}{\partial t} + \frac{y_i}{t} \frac{\partial \widehat{h}}{\partial y_i} \end{cases}$$

with the condition:

$$(9) \quad \frac{\partial \widehat{h}}{\partial z} = 0$$

This has the following consequences:

Remark 1.1. *Since $dt/ds = 0$, the flow preserves $t = \text{const.}$ Moreover*

$$L_{X_{\widehat{h}}}(\widehat{h}) = 0,$$

the flow in $\mathcal{M}$ associated to the vector field $X_{\widehat{h}}$ preserves the (constant) Hamiltonian function $\widehat{h}$.

Remark 1.2. For $t = 1$ and $\hat{h} = t \cdot h(x, y)$ one recovers the contact vector field defined in [3], that we call X_h^1 which is equal to:

$$(10) \quad \begin{cases} \frac{dx^i}{ds} = \frac{\partial h}{\partial y_i} \\ \frac{dy_i}{ds} = -\frac{\partial h}{\partial x^i} \\ \frac{dz}{ds} = -h + y_i \frac{\partial h}{\partial y_i} \end{cases}$$

Remark 1.3. The contactomorphisms generated by vector fields X_h^1 form a subgroup of the group of contact transformation. In fact, in [2] V. Arnol'd defined a contact Hamiltonian vector field without requiring that the associated one parameter transformation has the form (5). Arnol'd's definition leads to a vector field that generates a contact transformation but does not preserve the contact Hamiltonian $\hat{h}$.

Remark 1.4. Consider $\hat{h}_u : \mathcal{N} \rightarrow \mathbb{R}$ such that:

$$(11) \quad \hat{h}_u = \frac{1}{2}[-\langle \gamma x, x \rangle + 2t \langle \alpha x, y \rangle + t^2 \langle \beta x, y \rangle] + \langle v_1, y \rangle - \langle v_2, x \rangle \frac{1}{t} - kt$$

where α, β, γ are $n \times n$ real matrices with $\beta^T = \beta$ and $\gamma^T = \gamma$. One verifies that the vector field $X_{\hat{h}_u}$ at $t = 1$ is the infinitesimal generator of the one parameter group in $\mathbb{R}^{2n+2}$ $s \rightarrow \exp su$ with $u \in sp(2n+1, \mathbb{R})$, that is,

$$(12) \quad X_{\hat{h}_u} \Big|_{t=1} = \frac{d}{ds} \Big|_{s=0, t=1} \exp(su) \cdot (t, x, y, z).$$

Here

$$(13) \quad u = \begin{pmatrix} 0 & 0 & 0 \\ v & A & 0 \\ k & Jv & 0 \end{pmatrix},$$

where J is the standard symplectic structure in $\mathbb{R}^{2n}$, $v \in \mathbb{R}^{2n}$, $k \in \mathbb{R}$ and $A \in sp(2n, \mathbb{R})$ such that

$$(14) \quad JA = \begin{pmatrix} -\gamma & \alpha \\ \alpha^T & \beta \end{pmatrix},$$

Putting $(x, y) = w$, a simple computation shows that on level set $t = 1$:

$$(15) \quad X_{\hat{h}_u} \Big|_{t=1} = (v + Aw)^T \frac{\partial}{\partial w} + (k + (Jv)^T w) \frac{\partial}{\partial z}$$

2. THEORY OF CHARACTERISTICS FOR FIRST ORDER PDE AND ODD-SYMPLECTIC VECTOR FIELDS

In this section we shall follow the presentation of first order PDE given in [1]. Let us consider the case in which the manifold $\mathcal{M}$ is the first jet bundle associated to $\mathbb{R}^n$. Locally such space can be modeled by a $x \in \mathbb{R}^n$ and a germ of a function in x considered up to its gradient. To each $m \in J^1(\mathbb{R}^n)$ we associate $(x, z(x), \nabla z)$ with $z(\cdot)$ a germ of a smooth function in $\mathbb{R}^n$.

The manifold $J^1(\mathbb{R}^n)$ is a contact manifold with contact 1-form

$$\theta = dz - y_i dx^i$$

which vanishes on any germ $z(\cdot)$ such that $\partial z / \partial x^i = y_i$. The first jet space is the natural place to study the geometry of first order PDEs. In fact any PDE can be

understood as a submanifold of $J^1(R^n)$ (see [1]). Let us suppose that we have a smooth function on $J^1(R^n)$:

$$h : J^1(R^n) \rightarrow \mathbb{R} \text{ with } \frac{\partial h}{\partial z} = 0.$$

$h = 0$ defines a manifold in $J^1(R^n)$ which corresponds to the following PDE:

$$(16) \quad h(x, z(x), \nabla z(x)) = 0$$

To equation (16) we can associate a vector field X_c whose integral curves satisfy (10). We pose the natural question whether X_c is related to the *characteristic* vector field, which converts a first order PDE into a system of ordinary differential equations. In order to answer this question we need to recall the theory of characteristics. See [1] for more details.

2.1. Theory of characteristics for first order PDE. Let us assume that we have a first order PDE defined by the level set of $h : J^1(R^n) \rightarrow \mathbb{R}$. In the first jet space $J^1(\mathbb{R}^n)$ we can introduce *contact plane* Π_x at any $x \in \mathbb{R}^n$ spanned by the vectors V

$$(17) \quad i_V \theta = 0$$

Equation (17) determines a codimension 1 submanifold E^{2n} . Its tangent space TE^{2n} can be described by the vectors V such that:

$$(18) \quad i_V dh = 0$$

Definition 2.1. *The submanifold E^{2n} is called non-characteristic, if TE^{2n} is transversal to the contact plane Π .*

Definition 2.2. *The intersection of vector spaces*

$$P_x = T_x E^{2n} \cap \Pi_x$$

is called characteristic plane at x . Therefore P is defined by all vectors V such that

$$(19) \quad i_V \theta = 0 \text{ and } i_V dh = 0.$$

We shall consider only the case in which E^{2n} is not characteristic. Note that since TE^{2n} and Π lie in $J^1(R^n)$ we have:

$$\dim P_x = \dim TE_x^{2n} + \dim \Pi_x - \dim TJ^1(R^n) = 2n - 1$$

The 2-form $d\theta$, associated to the contact form θ , defines a skew product in $TJ^1(R^n)$. This allows us to select a vector in each characteristic plane P_x which is skew-orthogonal to P . The characteristic vector field for (16) is given by X_c :

- (1) $i_{X_c} \theta = 0$, X_c is a contact vector field,
- (2) $d\theta(X_c, V) = 0$ for all $V \in P$.

In symplectic geometry one can show that the skew-orthogonal complement to a k dimensional vector space is a $2n - k$ vector space. The characteristic plane P has dimension $2n - 1$. Therefore the characteristic direction X_c is uniquely determined. For a given $h : J^1(R^n) \rightarrow \mathbb{R}$, in local coordinates the vector field X_c has components:

$$(20) \quad \begin{cases} \frac{dx^i}{ds} = \frac{\partial h}{\partial y_i} \\ \frac{dy_i}{ds} = -\frac{\partial h}{\partial x^i} - y_i \frac{\partial h}{\partial z} \\ \frac{dz}{ds} = y_i \frac{\partial h}{\partial y_i} \end{cases}$$

2.2. Cauchy Problem. The theory of characteristics is used to solve the Cauchy problem. For an equation

$$h(x, z(x), \nabla z(x)) = 0$$

Cauchy data are: a manifold $\gamma \subset \mathbb{R}^n$ and the values $z|_\gamma = \phi(x)$. We have seen that the equation $h = 0$ corresponds to submanifold E^{2n} in $J^1(\mathbb{R}^n)$. Cauchy data define a submanifold $N \subset E^n$ called the *manifold of initial data*. It turns out that the Cauchy problem is solvable (locally) if N is not characteristic, that is if the projection of the characteristic direction to $\mathbb{R}^n$ is transversal to the tangent plane to N (see [1]). Given a parametrization $x_\gamma(\lambda)$ of γ , it is possible to construct the family of initial data:

$$x_0 = x_\gamma(\lambda), z_0 = \phi(x_\gamma(\lambda)), p_0 = p_0(\lambda)$$

where $p_0(\lambda)$ is such that:

$$h(x_\gamma(\lambda), \phi(x_\gamma(\lambda)), p_0(\lambda)) = 0$$

which forms the family of initial data for the characteristic vector field (20).

Remark 2.1. *The characteristic vector field (20) does not depend on whether the PDE is given by $h = 0$ or $h = c$ with $c \neq 0$, but the Cauchy problem does. In fact the value of the level set of h is contained in p_0 .*

3. THE MAIN RESULT

We can state:

Theorem 3.1. *For all the PDEs represented in $J^1(R^n)$ by a smooth function $h : J^1(R^n) \rightarrow \mathbb{R}$ of the form:*

$$h(x, \nabla z(x)) = 0,$$

the odd-symplectic Hamiltonian vector field (10) commutes with the characteristic vector field (20). Moreover on $h = 0$, the two vector fields coincide.

Proof. Consider the function $h : J^1(\mathbb{R}^n) \rightarrow \mathbb{R}$ as a contact Hamiltonian. Then by inspection one finds that the infinitesimal generator X_h^1 can be written as follows:

$$(21) \quad X_h^1 = X_c - h \frac{\partial}{\partial z}.$$

Since h does not depend on z , therefore $X_c(h) = 0$. Also the Lie bracket:

$$[X_c, X_h^1] = - \left[X_c, h \frac{\partial}{\partial z} \right] = 0$$

vanishes.

Now we want to compare X_h^1 (10) and X_c (20). We notice that if $h : J^1(R^n) \rightarrow \mathbb{R}$ does not depend on z , the first two components are equal. The component dz/ds differs because of the term $-h$ in (10). We have

$$L_{X_h^1}(h) = 0$$

that is h is preserved by the flow of X_h^1 . Therefore in the manifold $h = 0$ the vector field X_h^1 is equal to the characteristic vector field X_c $\square$

3.1. First order PDE and $sp(2n+1, \mathbb{R})$. Let us consider a particular class of first order PDEs in $\mathbb{R}^n$, namely,

$$(22) \quad c_{ij} \frac{\partial z(x)}{\partial x^i} \frac{\partial z(x)}{\partial x^j} + b_{ij} \frac{\partial z(x)}{\partial x^i} x^j + a_{ij} x^i x^j + e_i \frac{\partial z(x)}{\partial x^i} + f_i x^i = h_0$$

or more compactly:

$$(23) \quad \langle B \nabla z(x), \nabla z(x) \rangle + \langle e, \nabla z(x) \rangle + \langle f, x \rangle = h_0$$

Then we have:

Proposition 3.1. *For a given first order PDE of the form (23), if*

$$B \doteq \frac{1}{2} \begin{pmatrix} 2a & b \\ b^T & 2c \end{pmatrix}$$

(where $.^T$ is the transpose) is symmetric, then we can associate $u \in sp(2n+1, \mathbb{R})$

$$(24) \quad u = \begin{pmatrix} 0 & 0 & 0 \\ v & A & 0 \\ h_0 & Jv & 0 \end{pmatrix}$$

where $k = h_0$, $Jv = (e, f)$ and $-JB = A$, to the characteristic vector field of (22). If $a = b = 0$ and $v = (v_1, 0)$ in B then the characteristic vector field X_c associated to (22) commutes u .

Proof. The proof is essentially based on theorem 3.1 and on remark 1.4. According to theorem 3.1, equation (23) can be interpreted as a contact Hamiltonian $h(x, y)$ where $y = \nabla z(x)$, and the contact vector field on $h = 0$ is equal to the characteristic vector field. If the conditions on the matrices a, b, c are satisfied the PDE (23), on $J^1(\mathbb{R}^n)$ gives rise to a contact Hamiltonian of the form:

$$\widehat{h}(x, y, 1) = \langle w, JAw \rangle + \langle Jv, w \rangle + h_0, \text{ where } w = (x, y)$$

According to remark 1.4, we can associate to such Hamiltonian an element of $sp(2n+1, \mathbb{R})$ of the form (24).

Let us consider the Hamiltonian

$$(25) \quad \widehat{h}_u = \langle ax, x \rangle + 2t \langle bx, y \rangle + t^2 \langle cx, y \rangle + \langle v_1, y \rangle - \langle v_2, x \rangle \frac{1}{t} - h_0 t$$

The corresponding infinitesimal generator $X_{\widehat{h}_u}^1$ is:

$$(26) \quad X_{\widehat{h}_u}^1 = \xi + Z_{\widehat{h}} \frac{\partial}{\partial z},$$

where

$$(27) \quad Z_{\widehat{h}} = \langle v_1, y \rangle - \langle v_2, x \rangle + h_0$$

The characteristic vector field X_c of (23) is

$$(28) \quad X_c = \xi + Z_c \frac{\partial}{\partial z},$$

where

$$(29) \quad \xi = 2(bx + cy) \cdot \frac{\partial}{\partial x} - 2(ax + by) \cdot \frac{\partial}{\partial y}$$

$$(30) \quad Z_c = 2 \langle bx, y \rangle + 2 \langle cy, y \rangle + \langle v_1, y \rangle$$

Since h does not depend on z , the commutator is given by:

$$[X_{\hat{h}}^1, X_c] = (\xi(Z_c) - \xi(Z_{\hat{h}})) \frac{\partial}{\partial z}$$

After some algebra one finds:

$$(31) \quad \begin{aligned} \xi(Z_c) - \xi(Z_{\hat{h}}) = & 4 \langle bx, b^T y \rangle + 2 \langle cy, b^T y \rangle + \langle bx, v_2 \rangle + \langle cy, v_2 \rangle + \\ & -2 \langle bx, ax \rangle - 2 \langle bx, by \rangle - 4 \langle ax, cy \rangle - 4 \langle by, cy \rangle \end{aligned}$$

Last expression is identically zero for every (x, y) only for $a = b = 0$ and $v_2 = 0$. $\square$

3.2. Hamilton-Jacobi theory. In certain cases in Hamiltonian mechanics the construction of coordinate transformations that allows to integrate explicitly the equations of motion is based on the solution of Hamilton-Jacobi equation. We refer the reader to [1]. Suppose that we have a Hamiltonian system on $(\mathbb{R}^{2n}, \omega)$. The Cartan one-form, written in local coordinates, is:

$$(32) \quad \Theta = p_i dq^i - H(p, q, t) dt$$

where $H : \mathbb{R}^{2n} \times \mathbb{R} \rightarrow \mathbb{R}$ is the Hamiltonian. Θ is an integral invariant for the Hamiltonian flow. Consider the extended phase space $\mathcal{M} = \mathbb{R}^{2n} \times \mathbb{R}$ and two set of coordinates:

$$(q, p, t) \text{ and } (x, y, \tau)$$

There are two functions $K(x, y, \tau)$ and S such that

$$(33) \quad p_i dq^i - H(p, q, t) dt = y_i dx^i - K(x, y, \tau) d\tau + dS$$

The function S is called a “generating function” because it allows one to define a canonical transformation.

Equation (33) can be used to find the canonical transformation which realizes $(q, p, t) \rightarrow (x, y, \tau)$. This transformation is usually consider in the case $t = \tau$. In this case, by means of Legendre transformation one can write (33) as:

$$(34) \quad p_i dq^i - H(p, q, t) dt = x^i dy_i - K(x, y, t) dt + dS(q, y, t).$$

Then one usually chooses $K = 0$ and obtains:

$$(35) \quad \begin{cases} \frac{\partial S}{\partial y_i} = x_i \\ \frac{\partial S}{\partial q^i} = p_i \\ H\left(q, \frac{\partial S}{\partial q}\right) + \frac{\partial S}{\partial t} = 0 \end{cases}$$

The third equation of (35) is the Hamilton-Jacobi equation and determines the function S which completely describes the canonical transformation $(q, p, t) \rightarrow (x, y, t)$. The Hamilton-Jacobi equation is defined in $J^1(\mathbb{R}^{n+1})$ and fulfills the conditions of theorem 3.1.

3.3. Eikonal equation. In geometric optics and WKB approximation of quantum mechanics WKB approximation the Eikonal equation is relevant. This is a first order PDE given by:

$$(36) \quad \frac{1}{2} \delta^{ij} \frac{\partial S}{\partial x^i} \frac{\partial S}{\partial x^j} - N(x) = 0$$

S represents the wave front and N is a function:

$$N(x) = (n(x)/c)^2,$$

where in optics is square ratio of the refraction index $n(x)$ to the speed of light c ; where in WKB theory:

$$N(x) = V(x) - E,$$

namely, potential minus the in energy.

When in optics we consider a medium with a piece-wise constant refraction index (multilayers), $N(x)$ can be easily approximated by piece-wise constant function. In quantum mechanics this happens when the potential is piece-wise constant. In these cases proposition 3.1 can be applied in such layer. For instance if

$$N(x) = n_k^2/c^2 \text{ for } x \text{ in the } k\text{-layer}$$

then in k^{th} -layer we have symmetry generated by

$$(37) \quad u = \begin{pmatrix} 0 & 0 & 0 \\ 0 & -JA & 0 \\ n_k^2/c^2 & 0 & 0 \end{pmatrix}$$

with

$$G = \frac{1}{2} \begin{pmatrix} 0 & 0 \\ 0 & \mathbf{1} \end{pmatrix}.$$

A similar analysis can be carried out in the case of semi-classical description of system which have piece-wise constant potential e.g. multi-well potential.

ACKNOWLEDGMENTS

I really wish to thank Richard Cushman for his useful suggestions and long discussions we had at the Warwick University Mathematics Institute and his careful reading my manuscript. I want also to thank J.Sniatycki, and L.Bates for the interesting and stimulating discussions.

This paper has been supported by Marie Curie Fellowship (Contract n. HPMFCT-2000-00541) and by M.A.S.I.E. European network.

REFERENCES

- [1] Arnol'd V I 1988 *Geometrical methods in the theory of ordinary differential equations* Springer Verlag.
- [2] Arnol'd V I 1989 *Mathematical methods in Classical Mechanics* Springer Verlag.
- [3] Bates L, R Cushman 2002 *The Odd symplectic group in geometry* Preprint, Calgary
- [4] Workshop on "Geometry of momentum map and Hamiltonian Mechanics" (UMIST - Manchester July 2002)
- [5] Cushman R, Westbury B 2002 *Conjugacy classes in the odd symplectic group* Preprint, Warwick
- [6] Gelfand I M, Zelevinsky A V, 1984 *Models of representations and their hidden symmetries* Akad. Nauk SSSR, Insti. Prikl. Mat. **71** 26pp
- [7] Proctor A R, 1988 *Odd symplectic groups* Inventiones Math. **92** 307-332
- [8] McDuff D, Salamon D 1998 *Introduction to symplectic topology* Oxford Uni.Press

MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, CV4 7AL COVENTRY, U.K., **e-mail:** SBANO@MATHS.WARWICK.AC.UK, L.SBANO@MCLINK.IT